



Patient name: Jane Doe DOB:	Sample type: Blood Sample collection date:	Report date: Invitae #:
Sex: Female	Sample collection date:	Clinical team:
MRN:		

#### Reason for testing

Family history

#### Test performed

Sequence analysis and deletion/duplication testing of the 47 genes listed in the Genes Analyzed section.

Invitae Common Hereditary Cancers Panel



### **RESULT: NEGATIVE**

#### About this test

This diagnostic test evaluates 47 gene(s) for variants (genetic changes) that are associated with genetic disorders. Diagnostic genetic testing, when combined with family history and other medical results, may provide information to clarify individual risk, support a clinical diagnosis, and assist with the development of a personalized treatment and management strategy.

#### Next steps

- This test did not identify any pathogenic variants known to cause disease. This result should be discussed with a healthcare provider, such as a genetic counselor, to learn about the appropriate next steps for further evaluation. Clinical follow up may still be warranted. This result should be interpreted within the context of additional laboratory results, family history and clinical findings.
- Register your test at www.invitae.com/patients to download a digital copy of your results. You can also access educational resources about how your results can help inform your health.

### **Clinical Summary**

No reportable genetic variants were identified by this analysis, however, this individual may still be at risk for certain medical conditions based on other factors such as family history, genetic causes not evaluated with this test, or other environmental influences. Follow up of this individual and surveillance of their family members may still be indicated.



Patient name: Jane Doe

# **Genes Analyzed**

This table represents a complete list of genes analyzed for this individual, including the relevant gene transcript(s). If more than one transcript is listed for a single gene, variants were reported using the first transcript listed unless otherwise indicated in the report. Results are negative unless otherwise indicated in the report. Benign and Likely Benign variants are not included in this report but are available upon request. An asterisk (\*) indicates that this gene has a limitation. Please see the Limitations section for details.

GENE	TRANSCRIPT
APC	NM_000038.5
ATM	NM_000051.3
AXIN2	NM_004655.3
BARD1	NM_000465.3
BMPR1A	NM_004329.2
BRCA1	NM_007294.3
BRCA2	NM_000059.3
BRIP1	NM_032043.2
CDH1	NM_004360.3
CDK4	NM_000075.3
CDKN2A (p14ARF)	NM_058195.3
CDKN2A (p16INK4a)	NM_000077.4
CHEK2	NM_007194.3
CTNNA1	NM_001903.3
DICER1	NM_177438.2
EPCAM*	NM_002354.2
GREM1*	NM_013372.6
HOXB13	NM_006361.5
КІТ	NM_000222.2
MEN1	NM_130799.2
MLH1	NM_000249.3
MSH2	NM_000251.2
MSH3	NM_002439.4
MSH6	NM_000179.2
MUTYH	NM_001128425.1
NBN	NM_002485.4
NF1	NM_000267.3
NTHL1	NM_002528.6
PALB2	NM_024675.3
PDGFRA	NM_006206.4
PMS2	NM_000535.5
POLD1	NM_002691.3
POLE	NM_006231.3
PTEN	NM_000314.4
RAD50	NM_005732.3

GENE	TRANSCRIPT
RAD51C	NM_058216.2
RAD51D	NM_002878.3
SDHA*	NM_004168.3
SDHB	NM_003000.2
SDHC	NM_003001.3
SDHD	NM_003002.3
SMAD4	NM_005359.5
SMARCA4	NM_001128849.1
STK11	NM_000455.4
TP53	NM_000546.5
TSC1	NM_000368.4
TSC2	NM_000548.3
VHL	NM_000551.3



Patient name: Jane Doe

## Methods

- Genomic DNA obtained from the submitted sample is enriched for targeted regions using a hybridization-based protocol, and sequenced using Illumina technology. Unless otherwise indicated, all targeted regions are sequenced with ≥50x depth or are supplemented with additional analysis. Reads are aligned to a reference sequence (GRCh37), and sequence changes are identified and interpreted in the context of a single clinically relevant transcript, indicated below. Enrichment and analysis focus on the coding sequence of the indicated transcripts, 10bp of flanking intronic sequence (20bp for BRCA1/2), and other specific genomic regions demonstrated to be causative of disease at the time of assay design. Promoters, untranslated regions, and other non-coding regions are not otherwise interrogated. For some genes only targeted loci are analyzed (indicated in the table above). Exonic deletions and duplications are called using an in-house algorithm that determines copy number at each target by comparing the read depth for each target in the proband sequence with both mean read-depth and read-depth distribution, obtained from a set of clinical samples. Markers across the X and Y chromosomes are analyzed for quality control purposes and may detect deviations from the expected sex chromosome complement. Such deviations may be included in the report in accordance with internal guidelines. All clinically significant observations are confirmed by orthogonal technologies, except individually validated variants and variants previously confirmed in a first-degree relative. Confirmation technologies include any of the following: Sanger sequencing, Pacific Biosciences SMRT sequencing, MLPA, MLPA-seq, Array CGH. Array CGH confirmation of NGS CNV calling performed by Invitae Corporation (1400 16th Street, San Francisco, CA 94103, #05D2040778). The following analyses are performed if relevant to the requisition. For PMS2 exons 12-15, the reference genome has been modified to force all sequence reads derived from PMS2 and the PMS2CL pseudogene to align to PMS2, and variant calling algorithms are modified to support an expectation of 4 alleles. If a rare SNP or indel variant is identified by this method, both PMS2 and the PMS2CL pseudogene are amplified by long-range PCR and the location of the variant is determined by Pacific Biosciences (PacBio) SMRT sequencing of the relevant exon in both long-range amplicons. If a CNV is identified, MLPA or MLPA-seq is run to confirm the variant. If confirmed, both PMS2 and PMS2CL are amplified by long-range PCR, and the identity of the fixed differences between PMS2 and PMS2CL are sequenced by PacBio from the long-range amplicon to disambiguate the location of the CNV. Technical component of confirmatory sequencing is performed by Invitae Corporation (1400 16th Street, San Francisco, CA 94103, #05D2040778). Technical component of Fibroblast cell-culturing and gDNA extraction from skin punch biopsy is performed by Invitae Corporation (310 Goddard, Suite 150, Irvine CA 92618, #05D1052995).
- A PMID is a unique identifier referring to a published, scientific paper. Search by PMID at http://www.ncbi.nlm.nih.gov/pubmed.
- An rsID is a unique identifier referring to a single genomic position, and is used to associate population frequency information with sequence changes at that position. Reported population frequencies are derived from a number of public sites that aggregate data from large-scale population sequencing projects, including ExAC (http://exac.broadinstitute.org), gnomAD (http://gnomad.broadinstitute.org), and dbSNP (http://ncbi.nlm.nih.gov/SNP).
- A MedGen ID is a unique identifier referring to an article in MedGen, NCBI's centralized database of information about genetic disorders and phenotypes. Search by MedGen ID at http://www.ncbi.nlm.nih.gov/medgen. An OMIM number is a unique identifier referring to a comprehensive entry in Online Mendelian Inheritance of Man (OMIM). Search by OMIM number at http://omim.org/.
- Invitae uses information from individuals undergoing testing to inform variant interpretation. If "Invitae" is cited as a reference in the variant details this may refer to the individual in this requisition and/or historical internal observations.

### Limitations

Based on validation study results, this assay achieves >99% analytical sensitivity and specificity for single nucleotide variants, insertions and deletions <15bp in length, and exon-level deletions and duplications. Invitae's methods also detect insertions and deletions larger than 15bp but smaller than a full exon but sensitivity for these may be marginally reduced. Invitae's deletion/duplication analysis determines copy number at a single exon resolution at virtually all targeted exons. However, in rare situations, single-exon copy number events may not be analyzed due to inherent sequence properties or isolated reduction in data quality. Certain types of variants, such as structural rearrangements (e.g. inversions, gene conversion events, translocations, etc.) or variants embedded in sequence with complex architecture (e.g. short tandem repeats or segmental duplications), may not be detected. Additionally, it may not be possible to fully resolve certain details about variants, such as mosaicism, phasing, or mapping ambiguity. Unless explicitly guaranteed, sequence changes in the promoter, non-coding exons, and other non-coding regions are not covered by this assay. Please consult the test definition on our website for details regarding regions or types of variants that are covered or excluded for this test. This report reflects the analysis of an



Patient name: Jane Doe

extracted genomic DNA sample. In very rare cases (such as circulating hematolymphoid neoplasm, bone marrow transplant, recent blood transfusion, or maternal cell contamination), the analyzed DNA may not represent the patient's constitutional genome.

EPCAM: Sequencing analysis is not offered for this gene. SDHA: Deletion/duplication analysis is not offered for this gene. GREM1: Promoter region deletion/duplication testing only.

## Disclaimer

DNA studies do not constitute a definitive test for the selected condition(s) in all individuals. It should be realized that there are possible sources of error. Errors can result from trace contamination, rare technical errors, rare genetic variants that interfere with analysis, recent scientific developments, and alternative classification systems. This test should be one of many aspects used by the healthcare provider to help with a diagnosis and treatment plan, but it is not a diagnosis itself. This test was developed and its performance characteristics determined by Invitae. It has not been cleared or approved by the FDA. The laboratory is regulated under the Clinical Laboratory Improvement Act (CLIA) as qualified to perform high-complexity clinical tests (CLIA ID: 05D2040778). This test is used for clinical purposes. It should not be regarded as investigational or for research.

This report has been reviewed and approved by:

**Placeholder for signature**